


Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Asian Pacific Journal of Tropical Medicine

journal homepage: www.elsevier.com/locate/apjtm

Document heading doi:

Human cystic echinococcosis: epidemiologic, zoonotic, clinical, diagnostic and therapeutic aspects

Shyamapada Mandal^{1*}, Manisha Deb Mandal²¹Department of Zoology, Gurudas College, Narkeldanga, Kolkata–700 054, India²Department of Physiology and Biophysics, KPC Medical College and Hospital, Jadavpur, Kolkata–700 032, India

ARTICLE INFO

Article history:

Received 20 October 2011

Received in revised form 15 January 2012

Accepted 15 March 2012

Available online 20 April 2012

Keywords:

Human cystic echinococcosis

Zoonotic disease

Epidemiology

Echinococcus granulosus

Hydatid cyst

Surgery

Chemotherapy

ABSTRACT

This review represents an updated scenario on the transmission cycle, epidemiology, clinical features and pathogenicity, diagnosis and treatment, and prevention and control measures of a cestode parasite *Echinococcus granulosus* (*E. granulosus*) infection causing cystic echinococcosis (CE) in humans. Human CE is a serious life-threatening neglected zoonotic disease that occurs in both developing and developed countries, and is recognized as a major public health problem. The life cycle of *E. granulosus* involves a definitive host (dogs and other canids) for the adult *E. granulosus* that resides in the intestine, and an intermediate host (sheep and other herbivores) for the tissue-invading metacestode (larval) stage. Humans are only incidentally infected; since the completion of the life cycle of *E. granulosus* depends on carnivores feeding on herbivores bearing hydatid cysts with viable protoscoleces, humans represent usually the dead end for the parasite. On ingestion of *E. granulosus* eggs, hydatid cysts are formed mostly in liver and lungs, and occasionally in other organs of human body, which are considered as uncommon sites of localization of hydatid cysts. The diagnosis of extrahepatic echinococcal disease is more accurate today because of the availability of new imaging techniques, and the current treatments include surgery and percutaneous drainage, and chemotherapy (albendazole and mebendazole). But, the wild animals that involve in sylvatic cycle may overlap and interact with the domestic sheep–dog cycle, and thus complicating the control efforts. The updated facts and phenomena regarding human and animal CE presented herein are due to the web search of SCI and non-SCI journals.

1. Introduction

Echinococcosis, also called hydatid disease or hydatidosis, is a zoonosis, and in humans it occurs as a result of infection by the larval (metacestode) stages of taeniid cestodes of the genus *Echinococcus*. It is characterized by long-term growth of metacestode (larval) stages (hydatid cysts) in internal organs (mainly the liver and lungs) of intermediate host animals, viz., sheep and goats. Accidentally, eggs are also ingested by humans but do not play a role in the natural cycle. Six species of *Echinococcus* have been recognized; four are of public health concern, which are *Echinococcus granulosus* (*E. granulosus*) causing cystic echinococcosis (CE), *Echinococcus multilocularis* causing alveolar echinococcosis, and *Echinococcus vogeli* and *Echinococcus oligarthrus* causing polycystic echinococcosis; two new species have recently been identified: *Echinococcus*

shiquicus in small mammals from the Tibetan plateau^[1] and *Echinococcus felidis* in African lions^[2,3], but their zoonotic transmission potential is unknown. The *E. granulosus* infections have been further classified as domestic and sylvatic; in the domestic form, sheep serves as the most common intermediate host while in the sylvatic form the usual intermediate hosts are caribou or moose.

Human cystic echinococcosis is the most common presentation and probably accounts for > 95% of the estimated 2–3 million cases in the globe^[4]. Despite some progress in the control of echinococcosis, this zoonosis continues to be a major public health problem in several countries, while in many other countries the disease constitutes an emerging and re-emerging^[5]. Current analyses revealed that human CE is re-emerging problem with a remarkable economic impact in developed countries, viz., Spain^[6], and the disease continues to be a substantial cause of morbidity and mortality in several parts of the globe. The major sources of morbidity are pressure effects from cyst size, location in a sensitive organ (brain, reproductive tract, bone), or cyst rupture with subsequent anaphylaxis or dissemination of the infection.

*Corresponding author: Dr. Shyamapada Mandal, Department of Zoology, Gour Banga University, Mokdumpur, Malda–732103, India.
E-mail: samtropmed@gmail.com

The most common affected organs, however, are the liver and the lungs where 90% of the echinococcal cysts develop; excluding liver and lungs, all the other organs of the human body are considered as uncommon sites of localization of the hydatid disease. In this review, various aspects of human CE are discussed, based upon the current publications, in SCI and non-SCI journals, and in scientific websites, by different authors from different parts of the globe.

2. Etiologic identity

The adult *E. granulosus* (3–6 mm long) resides in the small intestine of the definitive hosts, dogs or other canids[5,7]. The adult worm consists of scolex bearing a rostellum with double row of numerous hooks, four prominent suckers and a short neck region, and the strobilae consist of 3 proglottids; usually one immature, one mature and one gravid proglottid make up the strobila.

The *E. granulosus* exists as a complex of different strains that differ in a wide variety of criteria having impact on the epidemiology and control of CE. The most important strain associated with the human CE appears to be the common sheep G1 strain, and is the most cosmopolitan form[5,8]. The current molecular characterization of human and animal *E. granulosus* isolates, however, demonstrated the involvement of the camel G6 strain in causing infection to humans[9,10].

3. Epidemiology

The greatest prevalence of the disease is in regions of the temperate countries, and this is an endemic disease in some parts of the world, such as South America, North Africa, Asia and Australia[11–13]. In the endemic parts, the rate of incidence of human CE has been recorded as >50 per 10⁵ person-years, and 5%–10% prevalence rate as in parts of Peru, Argentina, east Africa, central Asia, and China[4]; confirmed evidence for the emergence or re-emergence of human CE is there in parts of China, central Asia, Eastern Europe, and Israel[14]. The incidence of human hydatid cysts has been recorded as 18–20 per 10⁵ in Turkey, one of the endemic countries of the disease[15]. In Europe, human CE occurs in every country or region, where the annual incidence of hospital cases vary between <1 and >8 per 10⁵ population; in the northern part of Italy, the average annual CE incidence was recorded as 9.4–5.6/10⁵ inhabitants during 2003–2005[16]. However, Moro *et al*[17] reported that the actual prevalence of infection may greatly be underestimated by information from different official sources of the same geographical region; Figure 1 represents such phenomenon in Junin and in the coastal city of Chincha. As has been documented in literatures, human CE may occur in the subjects in between 1 and 75 year of ages[14]. Epidemiologically, CE occurs mostly in poor communities raising sheep and other livestock, and involving dogs in guarding as well as herding animals. *E. granulosus* is mainly transmitted in a cycle between dog definitive hosts and livestock (mainly sheep); the human behaviour also helps to perpetuate the domestic cycle of *E. granulosus*[4]. The CE is endemic in India; the foci of the disease exist in this

country with highest prevalence in Andhra Pradesh and Tamil Nadu[18]. Pednekar *et al*[19] reported, based upon the morphological and molecular studied, that *Echinococcus* of buffalo origin is phenotypically and genetically similar to the sheep (G1) and Tasmanian sheep (G2) strains of *Echinococcus* sensu stricto. In India, the annual incidence of human CE per 10⁵ persons varies from 1 to 200; it has been reported that human CE cases were with various clinical manifestations and immense morbidity[20].




Figure 1. Human cystic echinococcosis cases per 10⁵ from different sources of a particular geographic region.

4. Transmission cycle

The life cycles of *E. granulosus* can be classified as domestic, involving the domestic dog as the main definitive host and various species of domestic ungulates as intermediate hosts, or as sylvatic, involving wild carnivores (Foxes, wolves) and Cervidae (elk; *Alces alces*, reindeer; *Rangifer tarandus*, and red deer; *Cervus elaphus*) as hosts; the sylvatic cycle is reported in limited regions of the world such as North America and Eurasia[21]. In many areas of endemic infection, domestic and sylvatic life cycles coexist or overlap[7].

4.1. Natural cycle

The *E. granulosus* life cycle involves carnivores such as dogs (as well as wild canids like wolves and foxes) as definitive hosts in which the adult *E. granulosus* develops and resides in the small intestine; under natural conditions, herbivores mainly sheep (also goat, swine, cattle, horses and camels) serve as a suitable intermediate host where echinococcal cysts (larval stages) develop in various organs and tissues[5,14]. Humans function as accidental intermediate hosts, and they are usually the 'dead end' for the parasitic infection cycle; the consequences of human infection will be discussed later in this article.

Adult *E. granulosus* worms have been detected in various carnivores such as stray and farm dogs, red foxes, golden jackals and wolves[22–24]. Dogs play a critical role in transition the hydatidosis; the rate of infection with *E. granulosus* in stray dogs in Iran shows a prevalence of 5% – 49%[25]. The role of wild definitive hosts in the

epidemiology of *E. granulosus* is known[6]: prevalence of echinococcosis in wolves was found 14% in Spain, and the wolves parasitized with *E. granulosus* have also been found in Belarus, Italy, Finland, and Bulgaria (prevalence rate 11.5%–36%); all positive wolves harboured gravid *E. granulosus* from the sheep G1 strain. The wild definitive hosts may contribute in the transmission of animal (and human) CE in domestic cycles (Figure 2), since the G1 genotype has been found among wild animals.




Figure 2. Interaction between sylvatic cycle and domestic cycle of *E. granulosus* (indicated by bi-directional arrow) (A) In sylvatic cycle the definitive hosts acquire infection by scavenging herbivores acting as the intermediate hosts, (B) domestic cycle is maintained between dogs and sheep, with man as accidental intermediate host (not shown here).

4.2. Acquisition of infection by animals

Most forms of human CE are transmitted in domestic cycles involving dogs and livestock; the most known is the dog–sheep cycle. In a typical dog–sheep cycle, *E. granulosus* gravid proglottids or free eggs, which are passed in the faeces of infected dogs, are ingested by sheep (while grazing in the field contaminated with dog faeces), in which hydatid cysts (metacestode larval stages) develop[14]. The definitive host becomes infected by ingesting the cyst-containing organs of the infected intermediate host. Dogs usually acquire infection by ingestion of offal (liver and lungs) containing hydatid cysts with viable protoscoleces. Camels, which have an important role in transmission cycle of human CE, usually harbor G6 genotype of *E. granulosus*[21] that has also been reported from humans[26].

4.3. *E. granulosus* development

After ingestion by a suitable intermediate host (herbivore: sheep), the egg hatches in the small intestine and releases a hooked larva called oncosphere. The embryo, by means of its six hooks, penetrates the intestinal wall and migrates via the blood stream into major filtering organs, such as liver and lungs. In these organs, after localization, the oncosphere loses the hooks and develops into metacestode (larval echinococcal cyst or hydatid cyst) that enlarges gradually, producing many protoscoleces (future tapeworm heads) and daughter cysts that fill the cyst interior. The cycle is completed if the metacestode with fertile protoscoleces are eaten (along with the cyst-containing organs of the slaughtered parasitized sheep) by a suitable carnivore[27]. The different forms of the hydatid cysts may be of fertile, sterile and calcified types; occurrence such cysts in different organs among various intermediate host animals are represented in Figure 3. After ingestion, the protoscoleces evaginate, attach to the intestinal mucosa, and develop into adult stages in 32 to 80 days. Sexual maturity of adult *E. granulosus* occurs in the small intestine after 4–5 weeks of ingestion of offal containing viable protoscoleces. Eggs or gravid proglottids are shed in the faeces, and thus a new cycle begins.




Figure 3. Characteristic of livestock hydatid cysts.

4.4. Prevalence of hydatidosis among herbivores

The incidence of human CE is closely related to the prevalence of the disease in domestic herbivores, and currently there are reports on higher prevalence of bovine and ovine cystic hydatidosis[28]. The prevalence of infection with hydatid cyst in sheep, goats, cattle and buffaloes was recorded high in various endemic regions for *E. granulosus*[25,29]; involvement of herbivores as the intermediate hosts (bearing cystic hydatidosis) for the parasite is shown in Figure 4. Among the intermediate hosts, the most commonly infected organs are the lungs and liver[22,28,29]; the incidence of hydatid cysts in sheep, goat, cattle and camel and the organ involvement are shown in

Figure 5.




Figure 4. Cystic hydatidosis in animals acting as the intermediate hosts of *E. granulosus*.




Figure 5. Occurrence of hydatid cysts in different organs of some herbivores.

Rahimi *et al*[29] reported that the sheep is highly sensitive to the *E. granulosus* sheep G1 strain, and in this intermediate host the hydatid cysts are mostly fertile and, hence, the infected sheep are potential source of infection to dogs (and other carnivores), the reservoirs for human and livestock infections; most of the cysts were without protoscolices found in cattle having no important role in disease transmission. However, in Romania, as has been reported by Morariu *et al*[30], the livestock involved in the transmission pattern includes: sheep, cattle and swine. The prevalence of ovine cystic hydatidosis was recorded as 82.3% at the beginning of the control program in 1986, while in the year 2000 it was reduced to 20%[31]. Similar phenomena were seen for the other intermediate hosts[30]; the maximum and minimum occurrence of the disease has been depicted in the Figure 6.




Figure 6. Decreasing prevalence of cystic hydatidosis in herbivores.

5. Human infection (zoonotic aspect) of hydatidosis

5.1. Mode of infection

Humans become exposed to the eggs of the tapeworm after close contact with an infected dog or its contaminated environment[4]. The infected dogs pass in their feces *E. granulosus* eggs that adhere to hairs on the dog, and humans become exposed to the eggs after close contact with the dog (or its contaminated environment), and (humans) infected following accidental ingestion of *E. granulosus* eggs, or children in the course of playful and intimate contact with the infected dogs. Indirect transfer of *E. granulosus* eggs in contaminated water and uncooked food can also cause human infection. Certain human activities (*e.g.*, the widespread rural practice of feeding dogs the viscera of home-butchered sheep) facilitate transmission of the sheep strain and consequently increase the risk of human infection[32].

5.2. Development of cyst

In humans, as in the herbivorous intermediate hosts, hydatid cysts of *E. granulosus* develop mainly the liver and lungs as unilocular fluid-filled bladders, but the cysts may develop in almost any internal organ or tissue (heart, bone, muscle, nervous system) by haematogenous dissemination, or in the abdominal cavity[4,15]. It has been documented that most primary infections in humans result a single hydatid cyst; however, 20%–40% cases are reported to have multiple cysts or multiple organ involvement, which in the same patient may indicate ingestion of many eggs of *E. granulosus*[20]. Salem *et al*[27] described human hydatid cysts due to *E. granulosus* infection as fertile, sterile and calcified; Figure 7 depicts the occurrence of different forms of the cysts in human liver and lungs. The protoscolices are already formed in the fertile cysts, but a proportion of cysts does not produce protoscolices and remain sterile.




Figure 7. Characteristic of human hydatid cysts.

As has been recorded in the literatures, the cystic larva develops mainly in the liver (70%), in the lungs (20%), and 10% of cysts in anywhere of the body[7,33,34]; however, Fagzel and Ghanbary[35] documented that the liver is the most common site for echinococcal cyst development (>

65%), followed by the lungs (25%); the cyst occurs less frequently in the spleen and kidneys (3%), and heart, bone and brain (1%–2%). The cerebral hydatid cysts (comprising 1%–2%)[36,37] are classified as primary and secondary: primary cyst results from direct infestation of brain by the larva without any demonstrable involvement of other organs like liver and lungs[38]. Primary cysts are fertile containing brood capsules and scolices and their rupture may result in recurrent intracranial hydatid cyst; secondary cysts result from rupture of hydatid cyst in brain, and are infertile with negligible risk of recurrence after their rupture[39]. The hydatid cysts, which develop in bones in < 1% cases, were found as the most debilitating manifestation of *E. granulosus* infection to humans[40]. The proportion of hydatid cysts that occurs in muscle has been reported as 3%–5%; cysts in the left gluteus medius measure 4.8–5.6 cm in diameter, while the cyst developed in the gastrocnemius muscle was of 10 cm as has been detected by US and CT[41].

Larval infection (hydatid disease; hydatidosis) is characterised by long-term growth of metacestode (hydatid) cysts in humans. The diameters of cysts that generally increase are variable, 1–5 cm each year. Most *E. granulosus* cysts are 1–15 cm in diameter when they are discovered, but some may eventually reach 20 cm[7]. The cyst in a 50-year-old woman was of 18 cm, with daughter and granddaughter cysts, weighing about 1 kg, and three other cases with cyst size 10–18 cm (2 cases) and 20 cm (1 case) were females; human CE in pregnancy revealed that hydatid cysts may grow bigger due to decreased cell-mediated immunity and humoral effects of placental steroids, as has been seen in a study that the largest cyst (20 cm) was in the pregnant woman[20].

6. Clinical manifestation and pathogenesis

On ingestion of *E. granulosus* eggs by human (an accidental intermediate host), the embryos escape, penetrate the intestinal mucosa, enter the portal circulation, and are trapped in the liver; a small number escape the hepatic filter, enter the systemic circulation, and are scattered to other organs, where the oncospheres develop into fluid-filled unilocular (hepatic, or extra-hepatic) hydatid cysts.

The extra-hepatic hydatid cyst disease usually remains asymptomatic unless the cyst grows and produces symptoms due to pressure, rupture to the pleural or peritoneal cavity, secondary infection, or an allergic reaction. Before the slow growing echinococcal cysts enlarge sufficiently to cause symptoms, 5–20 years elapse; abdominal pain, hepatomegaly are the most common symptoms for patients with liver echinococcosis[42]. Because the larvae of this organism usually develop as discrete single cysts, it is the least severe and most treatable form; nevertheless, large or multiple cysts may cause irreversible damage to organs, and the rupture or puncture of the cyst can seed multiple organs with larvae or cause anaphylactic reactions. Cystic hydatidosis of the bone is often asymptomatic for a long duration and is usually detected only after a sudden fracture, secondary infection or neurovascular lesion caused by compression. Hemachander

et al[20] documented presenting complaints of CE cases including pain in abdomen, mass per abdomen, loss of appetite and obstructive jaundice.

The pre-operative complications include rupture of a cyst into a bronchus, bacterial infection of the cyst and obstructive jaundice, and the major complications after surgical resection with hepatic disease include single cases of pleural effusion, wound infection, biliary leak, and intra-abdominal bleeding resulting in a hepatic lobectomy[43]. Headache and vomiting due to raised intracranial tension are the most common presenting features of cerebral hydatid cysts; other manifestations include seizures, visual disturbances and ataxia[37,38].

7. Diagnosis


The useful diagnostic methods for human CE include ultrasonography (US), computerized tomography (CT) and magnetic resonance imaging (MRI)[41]. The US is the method of choice for the detection of both hepatic and extrahepatic echinococcal cysts; the sensitivity of this technique ranges 93%–98%[44]. The CT confirms the diagnosis by detecting the presence of daughter cysts and calcification in the cyst-wall; CT sensitivity ranges 90%–97%[44]. The MRI adequately demonstrates the features of the hydatid cyst; US and CT also contribute to pre-surgical diagnosis. The imaging with CT (as well as MRI) is central to the diagnosis of cerebral hydatid cysts[37]. Contrast enhanced CT scan of thorax is the diagnostic modality of choice for pulmonary hydatid cyst, and the “air bubble” sign (a relatively newly recognized radiological sign) has been reported to be very sensitive (85.7%) and specific (96.6%) in establishing diagnosis of ruptured, infected hydatid cyst[45,46]. The detection method of some extra hepato-pulmonary hydatid cysts along with their size and number and the actual localization, as has been reported by authors from different parts of the globe, are represented in Table 1.

The serological tests are helpful in making a human CE diagnosis. The current gold standard serodiagnosis is based on the detection of IgG antibodies to cyst-fluid derived native or recombinant antigen B subunits, either in ELISA or in immunoblots[4]. A novel 32 kDa EpC1 antigen, which is located in the germinal layer of the hydatid cyst and the early protoscolex of *E. granulosus*, demonstrated antibodies in 92.2% of preoperative human cases of CE compared with 84.5% cases detected using native antigen B[4]. Rahimi *et al*[29] assess the performance of antigen B (AgB) isolated from different *E. granulosus* intermediate hosts and from different cyst locations for immunodiagnosis of human CE, and reported that these increased the diagnostic sensitivity and specificity of the assay (Figure 8). Al-Sakee[47] showed that *E. granulosus* induces both cellular and humoral immune responses, and the number of peripheral blood CD8 T cells was significantly increased in cystic echinococcosis patients; this may help diagnosis of the disease.

Table 1

Occurrence of some extrahepato–pulmonary hydatid cysts in humans.

Localization	Size	Number	Detection	Country	Authors
Retroperitoneal	9 cm x4 cm	Single	US and CT	Greece	Tepetes <i>et al</i> [59]
Tibia	2.5–5 cm	Multiple	CT	Iran	Metanat <i>et al</i> [60]
Gastrocnemius muscle	10 cm	Double	CT	Turkey	Ormeci <i>et al</i> [41]
Gluteus medius muscle	< 5–5.6 cm	Multiple	US and CT	Turkey	Ormeci <i>et al</i> [41]
Gluteus maximus muscle	6 cm x3 cm	Single	CT	Turkey	Ormeci <i>et al</i> [41]
Superior rectus muscle	2 cm x2 cm	Single	MRI	Pakistan	Etaiwi and Mabreh[61]
Space in between right pelvis and right lumbar	17.5 cmx12.6 cm	Single	US	India	Roychowdhury <i>et al</i> [62]
Ovary	10 cm	Single	Surgical operation	Turkey	Vural <i>et al</i> [63]
Spleen, kidney, peritoneal cavity	4 cm x3 cm	Multiple	Laprotomy and US	India	Shukla <i>et al</i> [64]
Gallbladder	5 cm x3 cm	Single	Surgical operation	Morocco	Rabbani <i>et al</i> [65]
Vastuslateralis and rectus femoris muscles	17 cmx3 cmx5 cm	Single	CT	Hungary	Sisak <i>et al</i> [66]
Adductor magnus muscle of the left thigh	13 cmx8 cm	Single	MRI US	Italy	Vicidomini <i>et al</i> [67]
Right adductor brevis muscle	15 cmx15 cm	Single	MRI	Lebanon	George <i>et al</i> [68]
Left orbit	4 cmx1.8 cm	Single	CT and US	Tunisia	Limaïem <i>et al</i> [39]
Left thigh	10 cmx5 cmx4 cm	Single	CT, US	Greece	Mogrambi <i>et al</i> [69]

**Figure 8.** Diagnostic sensitivity and specificity of different AgB in diagnosis of human CE.

8. Treatment

The current methods of treatment of human CE include surgery (consisting of conservative and laparoscopic), and percutaneous drainage (consisting of puncture, aspiration, injection and re–aspiration; PAIR); these methods are mainly helpful for liver cysts. The method alternative to PAIR or surgery for the treatment of muscle–, spleen– or kidney–cysts include percutaneous drainage without re–aspiration[41,48,49]. The risk of anaphylactic reactions resulting from percutaneous treatment of echinococcal cysts has been found lower than that following administration of certain antibiotics, and fear of anaphylactic shock is no longer justified as an argument to avoid this therapeutic option[50].

Medical treatment with albendazole and mebendazole demonstrated efficacy is useful in the management of patients with hydatid cysts in liver and lungs. The combined chemotherapy with albendazole and praziquantel has been used successfully in the treatment of CE patients, and found more effective than monotherapy with these agents. Flubendazole alone, or in combination with ivermectin was reported effective in CE therapy[51]. Pre–surgical chemotherapy of *E. granulosus* infections reduces the size and number of viable protoscolices[52], and as has been documented the percutaneous treatment in combination with albendazole has been found more effective than

surgical cystectomy in terms of hospital stay (4.2 and 12.7 d, respectively) and complication rate (32% and 84%, respectively). The most widely used medications include mebendazole, albendazole, and praziquantel, but modifying the therapeutic approach by offering new pharmacological targets should be explored and evaluated[53,54].

9. Prevention and control

The human CE warrants more attention from clinicians, but their coordination with veterinarians and policy makers is also required to implement a more effective approach to control disease transmission Rojo–Vazquez *et al*[6]. Infection can be prevented through good personal hygiene to prevent hand–to–mouth transmission of *E. granulosus* eggs from dogs to humans, avoidance of ingestion of sheep viscera by dogs, by tailoring educational programmes, and by providing antihelminthic treatment to the definitive and intermediate animal hosts.

The current options for the control and prevention of hydatidosis include an effective livestock vaccine, potential dog vaccines against the worm *E. granulosus*, tailoring educational programmes, development of improved diagnostic tools for animal (definitive and intermediate) hosts and human beings, and more effective antiparasitic treatments like oxfendazole[4]. The biological cycle can be attacked at various points: regular dog deworming, controlled sheep slaughtering, vaccination of the intermediate (sheep) animal host, and in future vaccination of the definitive (dog) animal host[55–62]. However, since the wild animals involve in sylvatic cycle, which overlaps and interacts with the domestic sheep–dog cycle, this complicates the control efforts[6,14].

10. Concluding remarks

The infection of immature or mature intestinal stages of *E. granulosus* is not cause of morbidity to the definitive hosts, whereas the invasion of various organs (mainly liver and lungs) of humans and animal intermediate hosts by metacestodes can cause severe and fatal cystic hydatid disease. The disease is an important zoonosis, and is a serious public health and economic problem throughout the world.

It has been reported that the prevalence of hydatidosis in urban centres in India has been showing a consistently decreasing trend over the past few decades, which is possibly due to economic development and improved government legislation of abattoirs^[19]. A diagnosis of hepatic and extra-hepatic echinococcal disease is more accurate today because of the new imaging techniques (MRI, CT, US) available.

Study reinforces the importance of preoperative diagnosis with CT in preventing rupture of cyst and thus achieving optimal outcome as in case of cerebral hydatid cyst. En bloc resection without inducing rupture as well as spreading the daughter cyst is recommended treatment strategy, and accepted to be curative for intramuscular hydatid disease^[63,64].

The variation in *E. granulosus* may affect biology, host specificity and pathology, sensitivity to chemotherapeutics, and hence design and development of vaccines against *E. granulosus*, therefore, characterizing the exact etiological agent in different areas is imperative in order to determine the transmission patterns and control programs. Control programs based on combinations of animal vaccination, dehelminthization of dogs, and education programs should be implemented to achieve effective prevention of disease transmission^[65].

Conflict of interest statement

The authors declare no conflict of interest.

References

- [1] Xiao N, Qiu J, Nakao M, Yang W, Chen X, Schantz PM, et al. *Echinococcus shiquicus* n. sp., a taeniid cestode from Tibetan fox and plateau pika in China. *Int J Parasitol* 2005; **35**: 693–701.
- [2] Huttner M, Nakao M, Wassermann T, Siefert L, Boomker JD, Dinkel A, et al. Genetic characterization and phylogenetic position of *Echinococcus felidis* (Cestoda: Taeniidae) from the African lion. *Int J Parasitol* 2008; **38**: 861–868.
- [3] Huttner M, Siefert L, Mackenstedt U, Romig T. A survey of *Echinococcus* species in wild carnivores and livestock in East Africa. *Int J Parasitol* 2009; **39**: 1269–1276.
- [4] Craig PS, McManus DP, Lightowlers MW, Chabalgoity JA, Garcia HH, Gavidia CM, et al. Prevention and control of cystic echinococcosis. *Lancet Infect Dis* 2007; **7**: 385–394.
- [5] Moro P, Schantz PM. Echinococcosis: a review. *Int J Infect Dis* 2009; **13**: 125–133.
- [6] Rojo-Vazquez FA, Pardo-Lledias J, Francos-Von Hunefeld M, Cordero-Sanchez M, Alamo-Sanz R. Cystic echinococcosis in Spain: current situation and relevance for other endemic areas in Europe. *PLoS Negl Trop Dis* 2011; **5**: e893. doi:10.1371/journal.pntd.0000893
- [7] Ekert J, Deplazes P. Biological, epidemiological, and clinical aspects of echinococcosis, a zoonosis of increasing concern. *Clin Microbiol Rev* 2004; **17**: 107–135.
- [8] Wani MM, Durrani AM, Shafi M, Wani M, Khan M. Hydatid disease of the soft tissues of the lower limb: case report. *JK-Practitioner* 2007; **4**: 104–106.
- [9] Shahnazi M, Hejazi H, Salehi M, Reza Andalib AA. Molecular characterization of human and animal *Echinococcus granulosus* isolates in Isfahan, Iran. *Acta Trop* 2011; **117**: 47–50.
- [10] Magambo J, Njoroge E, Zeyhle E. Epidemiology and control of echinococcosis in sub-Saharan Africa. *Parasitol Int* 2006; **55**: 193–195.
- [11] Kilic D, Cangir AK, Bulut S, Akay H. Hydatid cyst of rib with intramedullary involvement. *Acta Chir Belg* 2003; **103**: 334–335.
- [12] Yang YR, Sun T, Li Z, Zhang J, Teng J, Liu X, et al. Community surveys and risk factor analysis of human alveolar and cystic echinococcosis in Ningxia Hui Autonomous Region, China. *Bull World Health Organ* 2006; **84**: 714–721.
- [13] Moro PL, Schantz PM. Echinococcosis: historical landmarks and progress in research and control. *Ann Trop Med Parasitol* 2006; **100**: 703–714.
- [14] McManus DP, Zhang W, Li J, Bartley PB. Echinococcosis. *Lancet* 2003; **362**: 1295–1304.
- [15] Kavukc S, Kilic D, Tokat AO, Kutlay H, Cangir AK, Enon S, et al. Parenchyma-preserving surgery in the management of pulmonary hydatid cysts: analysis of 1032 cases. *J Invest Surg* 2006; **19**: 60–68.
- [16] Logar J, Soba B, Kotar T. Serological evidence for human cystic echinococcosis in Slovenia. *BMC Infect Dis* 2008; **8**: 63. doi: 10.1186/1471-2334-8-63.
- [17] Moro PL, Budke CM, Schantz PM, Vasquez J, Santivanez SJ. Economic impact of cystic echinococcosis in Peru. *PLoS Negl Trop Dis* 2011; **5**: e1179. doi:10.1371/journal.pntd.0001179.
- [18] Nepalia S, Joshi A, Shende A, Sharma S. Management of echinococcosis. *J Assoc Physicians India* 2006; **54**: 458–462.
- [19] Pednekar RP, Gatne ML, Thompson RCA, Traub RJ. Molecular and morphological characterisation of *Echinococcus* from food producing animals in India. *Vet Parasitol* 2009; **165**: 58–65.
- [20] Hemachander SS, Prasad CR, Jessica M. Morbidity pattern of hydatid disease (cystic echinococcosis) and lack of its knowledge in patients attending mamata general hospital, Khammam, Andhra Pradesh. *Indian J Pathol Microbiol* 2008; **51**: 143–145.
- [21] Thompson RCA. The taxonomy, phylogeny and transmission of *Echinococcus*. *Exp Parasitol* 2008; **119**: 439–446.
- [22] Arbabi M, Hooshyar H. Survey of echinococcosis and hydatidosis in Kashan region, Central Iran. *Iranian J Publ Health* 2006; **35**: 75–81.
- [23] Sadjjadi SM. Present situation of echinococcosis in the Middle East and Arabic North Africa. *Parasitol Int* 2006; **55**: S197–S202.
- [24] Rokni MB. The present status of human helminthic diseases in Iran. *Ann Trop Med Parasitol* 2008; **102**: 283–295.
- [25] Rokni MB. Echinococcosis/hydatidosis in Iran. *Iranian J Parasitol* 2009; **4**: 1–16.
- [26] Manterola C, Benavente F, Melo A, Vial M, Roa JC. Description of *Echinococcus granulosus* genotypes in human hydatidosis in a region of southern Chile. *Parasitol Int* 2008; **57**: 342–346.
- [27] Salem CB, Schneegans F, Chollet JY, et Jemli MH. Epidemiological studies on echinococcosis and characterization of human and livestock hydatid cysts in Mauritania. *Iranian J Parasitol V* 2011; **6**: 49–57.
- [28] Fromsa A, Jobre Y. Infection prevalence of hydatidosis (*Echinococcus granulosus*, Batsch, 1786) in domestic animals in Ethiopia: A synthesis report of previous surveys. *Ethiop Vet J* 2011; **15**: 11–33.
- [29] Rahimi H, Sadjjadi SM, Sarkari B. Performance of antigen B isolated from different hosts and cyst locations in diagnosis of cystic echinococcosis. *Iranian J Parasitol* 2011; **6**: 12–19.
- [30] Morariu S, Bart JM, Cosoroaba M, Morariu F, Ilie M, Darabuş G. Prevalence of cystic echinococcosis in sheep, cattle and swine from the western Romania. *Sci Parasitol* 2011; **12**: 47–51.
- [31] Jimenez S, Perez A, Gil H, Schantz PM, Ramalle E, Juste RA. Progress in control of cystic echinococcosis in La Roja, Spain: decline in infection prevalences in human and animal hosts and economic costs and benefits. *Acta Trop* 2002; **83**: 213–221.
- [32] Moro PL, Caverio CA, Tambini M, Briceno Y, Jimenez R, Cabrera L. Identification of risk factors for cystic echinococcosis in a periurban population of Peru. *Trans Ryl Soc Trop Med Hyg* 2008; **102**: 75–78.
- [33] Jenkins DJ, Romig T, Thompson RC. Emergence/re-emergence of *Echinococcus* spp. – a global update. *Int J Parasitol* 2005; **35**: 1205–1209.
- [34] Rogan MT, Hai WY, Richardson R, Zeyhle E, Craig PS. Hydatid

- cysts: does every picture tell a story? *Trends Parasitol* 2006; **22**: 431–438.
- [35]Fagzel F, Ghanbary H. Hydatid cyst of the orbit. *J Isfahan Med School* 2002; **20**: 65.
- [36]Abdel Razek AA, El-Shamam O, Abdel Wahab N. Magnetic resonance appearance of cerebral cystic echinococcosis: World Health Organization (WHO) classification. *Acta Radiol* 2009; **50**: 549–554.
- [37]Wani NA, Kousar TL, Gojwari T, Robbani I, Singh M, zan AR, et al. Computed tomography findings in cerebral hydatid disease. *Turkish Neurosurg* 2011; **21**: 347–351.
- [38]Limaïem F, Bellil S, Bellil K, Chelly I, Mekni A, Khaldi M, et al. Primary hydatidosis of the central nervous system: a retrospective study of 39 Tunisian cases. *Clin Neurol Neurosurg* 2010; **112**: 23–28.
- [39]Limaïem F, Bellil S, Bellil K, Chelly I, Mekni A, Khaldi M, et al. Primary orbital hydatid cyst in an elderly patient. *Surg Infect* 2010; **11**: 393–395. doi:10.1089/sur.2009.019.
- [40]Zlitini M, Ezzaoula K, Lebib H, Karray M, Kooli M, Mesteri M. Hydatid cyst of bone: diagnosis and treatment. *World J Surg* 2001; **25**: 75–82.
- [41]Ormeci N, Idilman R, Akyar S, Palabiyikoglu M, Coban S, Erdem H, et al. Hydatid cysts in muscle: a modified percutaneous treatment approach. *Int J Infect Dis* 2007; **11**: 204–208.
- [42]Tsaroucha AK, Polychronidis AC, Lyrantzopoulos N, Pitiakoudis MS, Karayiannakis A, Manolas KJ, et al. Hydatid disease of the abdomen and other locations. *World J Surg* 2005; **29**: 1161–1165.
- [43]Somily A, Robinson JL, Miedzinski LJ, Bhargava R, Marrie TJ. Echinococcal disease in Alberta, Canada: more than a calcified opacity. *BMC Infect Dis* 2005; **5**: 34. doi:10.1186/1471-2334-5-34.
- [44]Balik AA, Celebi F, Basglu M, Oren D, Yildirgan I, Atamanalp SS. Intra-abdominal extrahepatic echinococcosis. *Surg Today* 2001; **31**: 881–884.
- [45]Yuncu G, Ors Kaya S, Sevinc S, Karabulut N, Alper H. The diagnostic value of the 'air bubble sign' in complicated pulmonary hydatid cysts. *J Thorac Cardiovasc Surg* 2007; **133**: 1524–1574.
- [46]Das A, Das SK. "Air-bubble" sign – An uncommon presentation of a common disease. *J Global Infect Dis* 2011; **3**: 310–312.
- [47]Al-Sakee HMA. Immunological aspects of cystic echinococcosis in ERBIL. *Zanco J Med Sci* 2011; **15**: 1–9.
- [48]Ormeci N, Soykan I, Palabiyikoglu M, Idilman R, Erdem H, Bektas A, et al. A new therapeutic approach for treatment of hydatid cysts of the spleen. *Dig Dis Sci* 2002; **47**: 2037–2044.
- [49]Ormeci N, Idilman R, Tuzun A, Erdem H, Palabiyikoglu M. A new percutaneous approach for the treatment of hydatid cyst of the kidney. *Int Urol Nephrol* 2005; **37**: 461–464.
- [50]Neumayr A, Troia G, de Bernardis C, Tamarozzi F, Goblirsch S, Piccoli L, et al. Justified concern or exaggerated fear: The risk of anaphylaxis in percutaneous treatment of cystic echinococcosis—A systematic literature review. *PLoS Negl Trop Dis* 2011; **5**: e1154. doi:10.1371/journal.pntd.0001154.
- [51]McManus DP. Echinococcosis with particular reference to Southeast Asia. In: D Rollinson, SI Hay, editors: *Advances in parasitology*, Vol. 72, Burlington: Academic Press; 2010, p. 267–303.
- [52]Miron D, Kavar B, Lumelsky D, Spiegel R, Horovitz Y. Encysted peritoneal hydatidosis in a child—the effectiveness of preoperative therapy with a combination of praziquantel and albendazole. *Ann Trop Med Parasitol* 2005; **99**: 577–582.
- [53]Voros D, Katsarelis D, Polymeneas G, Polydorou A, Pistiolis L, Kalovidouris A, et al. Treatment of hydatid liver disease. *Surg Infect (Larchmt)* 2007; **8**: 621–627.
- [54]Xiang-wei WU, Xue-ling C, Shi-jie Z, Xi Z, Hong S, Xin-yu P. Pericyst may be a new pharmacological and therapeutic target for hydatid disease. *Chinese Med J* 2011; **124**: 2857–2862.
- [55]Brunetti E, Garcia HH, Junghanss T. Cystic echinococcosis: chronic, complex, and still neglected. *PLoS Negl Trop Dis* 2011; **5**: e1146. doi:10.1371/journal.pntd.0001146.
- [56]Sathyanarayanan V, Sambhaji C, Saravu K, Razak A, Polnaya A, Rao SN. A rare case of hepatic cysticercosis. *Asian Pac J Trop Biomed* 2011; **1**(Suppl 1): S139–S140.
- [57]Bava AJ, Romero MM, Prieto R, Troncoso A. A case report of pulmonary coinfection of *Strongyloides stercoralis* and *Pneumocystis jiroveci*. *Asian Pac J Trop Biomed* 2011; **1**(4): 334–336.
- [58]Bava J, Lloveras S, Garro S, Troncoso A. Pulmonary coinfection by *Pneumocystis jiroveci* and *Cryptococcus neoformans*. *Asian Pac J Trop Biomed* 2011; **2**(1): 80–82.
- [59]Rakotoarivelo RA, Andrianasolo RL, Ranoharison D, Rakoto ROS, Sendrasoa FA, MJ de Dieu Randria. Neurocysticercosis as an important differential of paradoxical response during antituberculosis therapy in HIV–negative patient. *Asian Pac J Trop Dis* 2011; **1**(4): 333–334.
- [60]Gill M, Boombak E, Gill PS, Gupta V, Sen R. Extensive cardiac cysticercosis—an interesting autopsy finding. *Asian Pac J Trop Dis* 2011; **1**(4): 335–336.
- [61]Chakraborty A, Panja S, Bhattacharjee I, Chandra G, Hati A. A longitudinal study of neurocysticercosis through CT scan of the brain. *Asian Pac J Trop Dis* 2011; **1**: 206–208.
- [62]Wilma DSC, Kavya N, Kulkarni S. Evaluation of insulin sensitivity status in polycystic ovarian syndrome. *Asian Pac J Trop Dis* 2011; **1**: 67–70.
- [63]Parray FQ, Ahmad SZ, Sherwani AY, Chowdri NA, Wani KA. Primary paraspinous hydatid cyst: a rare presentation of echinococcosis. *Int J Surg* 2010; **8**: 404–406.
- [64]Arazi M, Erikoglu M, Odev K, Memik R, Ozdemir M. Primary echinococcus infestation of the bone and muscles. *Clin Orthopaed Related Res* 2005; **43**: 234–241.
- [65]Neghina R, Neghina AM, Marincu I, Iacobiciu I. Epidemiology and epizootology of cystic echinococcosis in Romania 1862–2007. *Foodborne Pathog Dis* 2010; **7**: 613–618.
- [66]Tepetes K, Christodoulidis G, Spyridakis M, Hatzitheoflou K. Large solitary retroperitoneal echinococcal cyst: A rare case report. *World J Gastroenterol* 2007; **13**: 6101–6103.
- [67]Metanat M, Sharifi–mood B, Sandoghi M, Alavi–Naini R. Osseous hydatid disease. *Iranian J Parasitol* 2008; **3**: 60–64.
- [68]Etaiwi M, Mabreh L. Intramuscular hydatid cyst of the superior rectus muscle. *Pak J Med Sci* 2008; **24**: 762–764.
- [69]Roychowdhury A, Bandopadhyay A, Bhattacharya P, Mitra RB. An unusual case of primary intrapelvic hydatid cyst. *Indian J Pathol Microbiol* 2010; **53**: 588–589.
- [70]Vural M, Yalcin S, Yildiz S, Camuzcuoglu H. Isolated type I pelvic cystic echinococcosis mimicking ovarian tumor. *North Am J Med Sci* 2011; **3**: 289–291.
- [71]Shukla S, Singh SK, Pujani M. Multiple disseminated abdominal hydatidosis presenting with gross hydatiduria: A rare case report. *Indian J Pathol Microbiol* 2009; **52**: 213–214.
- [72]Rabbani K, Narjis Y, Louzi A, Benelkhaïat R, Jalal H, Finech B. Unusual localization of hydatidosis: Hydatid cyst of gallbladder. *Ann Trop Med Public Health* 2011; **4**: 119–121.
- [73]Sisak CK, Bardocz L, Toth K. Pathological femoral neck fracture caused by an echinococcus cyst of the vastus lateralis—case report. *BMC Infect Dis* 2011; **11**: 103.
- [74]Vicidomini S, Cancerini G, Gabrielli S, Naspetti R, Bartoloni A. Muscular cystic hydatidosis: case report. *BMC Infect Dis* 2007; **7**: 23. doi:10.1186/1471-2334-7-23.
- [75]Abi Saad GS, Musallam KM, Korban ZR, Reslan OM, Mneimne M. Solitary hydatid cyst of the thigh: a challenging diagnosis. *Vector–Borne Zoonotic Dis* 2009; **9**: 743–745.
- [76]Mogrambi SAI, Fragandreas G, Papadopoulos S, Arapoglou S, Kambouridis A, Gerasimidis T. A rare case of femoral hydatidosis. *Surg Infect* 2010; **11**: 51–54.